

# The REDTOP experiment



*A. Mazzacane*



Rare Processes and Precision Frontier Townhall Meeting

October 2, 2020

# Rationale for an $\eta/\eta'$ Factory



- Recent results from the LHC suggest that the next search for New Physics should be performed in the low-energy mass range (MeV-GeV) using high-intensity beams.
- Light dark matter must be neutral under SM charges, otherwise it would have been discovered at previous colliders  
[G. Krnjaic RF6 Kickoff Meeting, August 12, 2020]
- The only known particles with all-zero quantum numbers are the  $\eta/\eta'$  mesons and the Higgs boson
- An  $\eta/\eta'$  factory is an excellent laboratory for precision measurements and a unique window to search for Physics Beyond the Standard Model in the MeV-GeV mass range.

# The REDTOP Experiment



- REDTOP is a proposed  $\eta/\eta'$  factory which aims at detecting small deviations from the Standard Model by collecting a large set of events from protons impinging on a fixed target
- The experiment will produce  $\sim 10^{13}$   $\eta$  mesons or  $\sim 10^{11}$   $\eta'$  mesons per year corresponding to an increase of the existing world sample by four order of magnitude
- All electromagnetic and strong decays of the neutral and long-lived  $\eta$  and  $\eta'$  are suppressed at first order and weak decays have branching ratios of order  $10^{-11}$
- Branching Ratio of processes from New Physics are enhanced compared to other systems.

# Physics Program



## C, T, CP-violation

- CP Violation via Dalitz plot mirror asymmetry:  $\eta \rightarrow \pi^0 \pi^+ \pi^-$
- CP Violation (Type I - P and T odd , C even):  $\eta \rightarrow 4\pi^0 \rightarrow 8\gamma$
- CP Violation (Type II - C and T odd , P even):  $\eta \rightarrow 4\pi^0 l^+ l^-$  and  $\eta \rightarrow 3\gamma$
- Test of CP invariance via  $\mu$  longitudinal polarization:  $\eta \rightarrow \mu^+ \mu^-$
- Test of CP invariance via  $\gamma^*$  polarization studies:  $\eta \rightarrow \pi^+ \pi^- e^+ e^-$  and  $\eta \rightarrow \pi^+ \pi^- \mu^+ \mu^-$
- Test of CP invariance in angular correlation studies:  $\eta \rightarrow \mu^+ \mu^- e^+ e^-$
- Test of T invariance via  $\mu$  transverse polarization:  $\eta \rightarrow \pi^0 \mu^+ \mu^-$  and  $\eta \rightarrow \gamma \mu^+ \mu^-$
- CPT violation:  $\mu$  polarization in  $\eta \rightarrow \pi^+ \mu^- \nu$  vs  $\eta \rightarrow \pi^- \mu^+ \nu$  and  $\gamma$  polarization in  $\eta \rightarrow \gamma \gamma$

## Other discrete symmetry violations

- Lepton Flavor Violation:  $\eta \rightarrow \mu^+ e^- + c.c.$
- Double lepton Flavor Violation:  $\eta \rightarrow \mu^+ \mu^+ e^- e^- + c.c.$

## Non- $\eta/\eta'$ based BSM Physics

- Dark photon and ALP searches in Drell-Yan processes:  $q\bar{q} \rightarrow A'/a \rightarrow l^+ l^-$
- $p + D \rightarrow {}^3\text{He}^+ + X_{17}$  with  $X_{17} \rightarrow e^+ e^-$
- ALP's searches in Primakoff processes:  $pZ \rightarrow pZa \rightarrow l^+ l^-$  (F. Kahlhoefer)
- Charged pion and kaon decays:  
 $\pi^+ \rightarrow \mu^+ \nu A' \rightarrow \mu^+ \nu e^+ e^-$  and  $K^+ \rightarrow \mu^+ \nu A' \rightarrow \mu^+ \nu e^+ e^-$
- Neutral pion decay:  $\pi^0 \rightarrow \gamma A' \rightarrow \gamma e^+ e^-$

## New particles and forces searches

- Scalar meson searches (charged channel):  $\eta \rightarrow \pi^0 H$  with  $H \rightarrow e^+ e^-$  and  $H \rightarrow \mu^+ \mu^-$
- Dark photon searches:  $\eta \rightarrow \gamma A'$  with  $A' \rightarrow l^+ l^-$
- QCD axion searches:  $\eta \rightarrow \pi^+ \pi^- a$  and  $\eta \rightarrow \pi^0 \pi^0 a$  with  $a \rightarrow l^+ l^-$
- Protophobic fifth force searches:  $\eta \rightarrow \gamma X_{17}$  with  $X_{17} \rightarrow e^+ e^-$
- New leptophobic baryonic force searches:  $\eta \rightarrow \gamma B$  with  $B \rightarrow e^+ e^-$  or  $B \rightarrow \gamma \pi^0$
- Indirect searches for dark photons new gauge bosons and leptoquark:  $\eta \rightarrow \mu^+ \mu^-$  and  $\eta \rightarrow e^+ e^-$
- Search for true muonium:  $\eta \rightarrow \gamma(\mu^+ \mu^-)|_{2M_\mu} \rightarrow \gamma e^+ e^-$

## Other Precision Physics measurements

- Proton radius anomaly:  $\eta \rightarrow \gamma \mu^+ \mu^-$  vs  $\eta \rightarrow \gamma e^+ e^-$
- All unseen leptonic decay mode of  $\eta/\eta'$  (SM predicts  $10^{-6} \div 10^{-9}$ )

## High precision studies on medium energy physics

- Nuclear models
- Chiral perturbation theory
- Non-perturbative QCD
- Isospin breaking due to the u-d quark mass difference
- Octet-singlet mixing angle
- Electromagnetic transition form-factors (important input for g-2)

- The physics sectors which can be probed at REDTOP range from the violation of discrete symmetries to the search for new particles  
[S. Tulin et al. <https://arxiv.org/abs/2007.00664>]
- Non-eta meson sectors can also be explored, such as ALPs and QCD-axions  
[D. S. M. Alves, arXiv:2009.05578]
- Finally, the Standard Model can be probed at low energy at an unprecedented precision level.

# $\eta$ Factories and Outstanding Anomalies



## Muonic puzzle (aka, proton radius and muon g – 2 Anomalies)

- Solved by postulating a new scalar boson  $\phi$  decaying into  $e^+e^-$ :

$$\eta \rightarrow \phi \pi^0 \rightarrow e^+e^- \gamma\gamma$$

$\eta \rightarrow \pi^0 e^- e^+$  is forbidden in the SM by charge conjugation symmetry at tree level but allowed by a virtual  $\phi$  emission

[Y. Liu,1, I.C. Cloët, G. A. Miller, *Nucl.Phys. B* (2019) 114638]

- Also interesting:

$$BR(\eta \rightarrow e^+e^-) \text{ vs } BR(\eta \rightarrow \mu^+\mu^-)$$

branching ratios (BR) in SM range from  $10^{-9}$  to  $10^{-6}$

[Pere Masjuan,a Pablo Sanchez-Puertas, *Phys.Rev.D* 26 (1982) 3302]

## X17 in the $e^+e^-$ emission spectra of isoscalar magnetic transitions of ${}^8Be$ and ${}^4He$ nuclei

- Solved by postulating a 17 MeV QCD-axion or a protophobic gauge boson decaying into  $e^+e^-$

$$\eta \rightarrow a \pi^+ \pi^- \rightarrow e^+e^- \pi^+ \pi^- \text{ and } \eta \rightarrow a \pi^0 \pi^0 \rightarrow e^+e^- \gamma\gamma\gamma\gamma$$

[D. S. M. Alves, arXiv:2009.05578]

$$\eta \rightarrow X17\gamma \rightarrow e^+e^- \gamma$$

[J.L. Feng et Al. *Phys. Rev. D* **95**, 035017]

## KOTO anomaly (100x excess of events in $K^0 \rightarrow \pi^0 \nu\nu$ )





- Solved by postulating a new Hadrophilic scalar boson  $H$  decaying into  $e^+e^-$

$$\eta \rightarrow H \pi^0 \rightarrow e^+e^- \gamma\gamma$$

[D. Egana-Ugrinovic , S. Homiller , and Patrick Meade, *Phys. Rev. Lett.* 124, 191801 (2020)]

[B. Batell, A. Freitas, A. Ismail, D. McKeen, *Phys. Rev. D* **100**, 095020]

# Sensitivity Studies at CERN PBC



Slide Credit: Susan Gardner & Jun Shi

University of Kentucky

# $\eta/\eta'$ Production Option-1: High Energy - Low Intensity Beam




- The production mechanism is based on the formation and the decay of intra-nuclear baryonic resonances
- $\eta$  and  $\eta'$  mesons are produced from a proton beam scattered on a target made of multiple thin Lithium (or Beryllium) foils
- Monte Carlo studies indicate optimal beam energies 1.8-2.1 GeV ( $\eta$ ) and 3.0-4.0 GeV ( $\eta'$ )
- Large production cross-section (several mbarn) relaxes the beam power requirement. A 30-50W beam is sufficient to generate the desired statistics.
- $\eta/\eta'$  mesons are produced with small boost and almost  $4\pi$  detector is necessary
- Hadronic inelastic background is reduced by using a Continuous Wave beam.
- Advantages:
  - Low intensity beam
- Disadvantages :
  - $\eta/\eta'$  are not tagged

# $\eta/\eta'$ Production Option-2: Low Energy - High Intensity Beam



- $\eta$  and  $\eta'$  are produced on a gaseous Deuterium target via the nuclear process



- A threshold of 880 MeV beam energy is required (1.7 GeV for  $\eta'$ )
- A larger beam intensity ( $\sim 1\text{MW}$ ) will compensate for a smaller production cross section  $\sim O(\mu\text{barn})$
- Advantages:
  - The combinatorics background from non- $\eta$  events is greatly reduced ( $\sim 10^8$  from preliminary studies) by tagging the  ${}^3\text{He}^{++}$  ion
  - The kinematics if fully closed and missing 4-P technique is applicable when searching for long lived particles escaping the detector same as B factories but with 40k x statistics
- Disadvantages:
  - A higher beam power is needed
  - An additional forward detector is required to tag the  ${}^3\text{He}^{++}$



# Experimental Techniques

- The interaction rate is large (1Ghz for option 1 and  $\sim$ 100Ghz for option 2) but event multiplicity is very low ( $\leq 8$  particles/event)
  - Crucial requirements are:
    - *Excellent PID for lepton/hadron separation*
    - *Fast timing with  $\sim$ 30ps resolution (for TOF purposes)*
- 
- Use multiple thin targets to minimize combinatorics background

## Charged tracks detection

### Option 1: Optical-TPC

Barions and most pions are below threshold

*Electrons and most muons are detected and reconstructed*

### Option 2: LGAD Tracker

4D track reconstruction for multihadron rejection

Complemented with quartz cells for TOF measurements

## $\gamma$ detection

### 5D-Calorimeter ADRIANO2

(PFA+Dual-readout)

### Sci and Cer light read by SiPM or SPAD

For excellent energy and position resolution and PID to disentangle showers from  $\gamma/\mu/\text{hadrons}$

## Fiber tracker (LHCb style)

for rejection of background from  $\gamma$ -conversion and reconstruction of secondary vertices  
( $\sim 70\mu\text{m}$  resolution)

# The REDTOP Detector



## Optical-TPC

For slow background rejection

or

## LGAD Tracker surrounded by Quartz cells

For 4D track reconstruction and TOF measurements



## 5D- Calorimeter: ADRIANO2 (Dual-readout +PFA)

Sci and Cer light read by SiPM or SPAD

For excellent energy, position resolution and PID

## $\mu$ -polarimeter (optional)

sandwich of fused silica and Si-pixel  
for measurement of muon polarization



## Vertex Fiber tracker

for rejection of  $\gamma$ -conversion and identifying displaced vertices from long lived particles

## 10x Be or Li targets

## Forward Detector for Option 2

for tagging  ${}^3\text{He}^{++}$  ions

# Collaboration

11 Countries, 35 Institutions, 92 Collaborators



J. Comfort, P. Mauskopf, D. McFarland, L. Thomas

Arizona State University, (USA)

I. Pedraza, D. Leon, S. Escobar, D. Herrera, D Silverio

Benemérita Universidad Autónoma de Puebla, (Mexico)

A. Alqahtani

Georgetown University, (USA)

F. Ignatov

Budker Institute of Nuclear Physics – Novosibirsk, (Russia)

A. Kotwal

Duke University, (USA)

M. Spannowsky

Durham University, (UK)

J. Dey, V. Di Benedetto, B. Dobrescu, E. Gianfelice-Wendt, E. Hahn, D. Jensen, C. Johnstone, J. Johnstone, J. Kilmer, G. Krnjaic, T. Kobilarcik, A. Kronfeld, K. Krempetz, M. May, A. Mazzacane, N. Mokhov, W. Pellico, A. Pla-Dalmau, V. Pronskikh, E. Ramberg, J. Rauch, L. Ristori, E. Schmidt, G. Sellberg, G. Tassotto, Y.D. Tsai

Fermi National Accelerator Laboratory, (USA)

P. Sanchez-Puertas

IFAE – Barcelona (Spain)

C. Gatto<sup>1†</sup>

Istituto Nazionale di Fisica Nucleare – Sezione di Napoli, (Italy)

W. Baldini

Istituto Nazionale di Fisica Nucleare – Sezione di Ferrara, (Italy)

R. Carosi, A. Kievsky, M. Viviani

Istituto Nazionale di Fisica Nucleare – Sezione di Pisa, (Italy)

W. Krzemień, M. Silarski, M. Zielinski

Jagiellonian University, Krakow, (Poland)

D. S. M. Alves, S. Pastore

Los Alamos National Laboratory, (USA)

M. Berlowski

National Centre for Nuclear Research – Warsaw, (Poland)

G. Blazey, A. Dychkant, K. Francis, M. Syphers, V. Zutshii, P. Chintalapati, T. Malla, M. Figora

Northern Illinois University, (USA)

D. Egaña-Ugrinovic

Perimeter Institute for Theoretical Physics – Waterloo, (Canada)

Y. Kahn

Princeton University – Princeton, (USA)

P. Meade, S. Homiller

Stony Brook University – New York, (USA)

A. Gutiérrez-Rodríguez, M. A. Hernandez-Ruiz

Universidad Autónoma de Zacatecas, (Mexico)

B. Fabela-Enriquez

Vanderbilt University, (USA)

J. Jaeckel

Universität Heidelberg, (Germany)

C. Siligardi, S. Barbi, C. Mugoni

Università di Modena e Reggio Emilia, (Italy)

L. E. Marcucci<sup>4</sup>

Universita' di Pisa, (Italy)

M. Guida<sup>3</sup>

Università di Salerno, (Italy)

S. Charlebois, J. F. Pratte

Université de Sherbrooke, (Canada)

L. Harland-Lang

University of Oxford, (UK)

S. Gori

University of California Santa Cruz, (USA)

R. Gardner, P. Paschos

University of Chicago, (USA)

J. Konisberg

University of Florida, (USA)

M. Murray, C. Rogan, C. Royon, Nicola Minafra, A. Novikov, F. Gautier, T. Isidori

University of Kansas, (USA)

S. Gardner, J. Shi, X. Yan

University of Kentucky, (USA)

M. Pospelov

University of Minnesota, (USA)

D. Gao

University of Science and Technology of China, (China)

K. Maamari

University of Southern California, (USA)

A. Kupsc

University of Uppsala, (Sweden)

S. Tulin

York University, (Canada)

# Summary



- The  $\eta/\eta'$  mesons provide an excellent laboratory to search for New Physics and precision tests of the Standard Model
- The REDTOP experiment is a  $\eta/\eta'$  factory designed to produce  $10^{13}$   $\eta$  and  $10^{11}$   $\eta'$  mesons
- The experiment is optimized to search for leptonic decays. It is fully complementary to JEF's searches
- Novel detector techniques need to be developed to cope with the high interaction rate
- Future High Energy and High Intensity experiments will benefit from the ensuing R&D.
- New studies are ongoing to improve the sensitivity for different running modes
- Multiple DOE laboratories have the capability of hosting REDTOP.